

Using The Binomial Distribution, If $N=7$ And $P=0.8$, Find $P(x=5)$. Round Your Answer To 4 Decimals.

Using The Binomial Distribution, If $N=7$ And $P=0.8$, Find $P(x=5)$. Round Your Answer To 4 Decimals.

Understanding the binomial distribution is essential for many statistical and probability applications, especially when dealing with scenarios involving a fixed number of independent trials, each with the same probability of success. In this article, we will explore how to calculate the probability $P(x=5)$ when the number of trials N is 7, and the probability of success in each trial P is 0.8. We will also discuss the binomial distribution formula in detail, illustrate the calculation step-by-step, and provide insights into interpreting the results.

What is the Binomial Distribution?

The binomial distribution is a discrete probability distribution that models the number of successes in a fixed number of independent Bernoulli trials. Each trial has only two possible outcomes—success or failure—and the probability of success remains constant throughout the trials.

Key Characteristics of the Binomial Distribution

- Number of trials: N (fixed in advance)
- Probability of success in each trial: P
- Probability of failure in each trial: $1 - P$
- Number of successes: x , where $x = 0, 1, 2, \dots, N$

The probability of observing exactly x successes in N trials is given by the binomial probability formula:

$$P(X = x) = \binom{N}{x} \times P^x \times (1 - P)^{N - x}$$

where $\binom{N}{x}$ is the binomial coefficient, calculated as:

$$\binom{N}{x} = \frac{N!}{x!(N-x)!}$$

$$\binom{N}{x} = \frac{N!}{x!(N-x)!}$$

\]

Applying the Binomial Formula to Our Problem

Given:

- Number of trials, $(N = 7)$
- Probability of success per trial, $(P = 0.8)$
- Desired number of successes, $(x = 5)$

Our goal is to compute:

\[

$$P(x=5) = \binom{7}{5} \times 0.8^5 \times (1 - 0.8)^{7-5}$$

\]

which simplifies to:

\[

$$P(x=5) = \binom{7}{5} \times 0.8^5 \times 0.2^2$$

\]

Step-by-Step Calculation

Let's proceed through the calculation systematically.

Step 1: Calculate the binomial coefficient $(\binom{7}{5})$

\[

$$\binom{7}{5} = \frac{7!}{5! \times (7-5)!} = \frac{7!}{5! \times 2!}$$

\]

Calculating factorials:

- $(7! = 7 \times 6 \times 5 \times 4 \times 3 \times 2 \times 1 = 5040)$
- $(5! = 5 \times 4 \times 3 \times 2 \times 1 = 120)$
- $(2! = 2 \times 1 = 2)$

Plugging in:

\[

$$\binom{7}{5} = \frac{5040}{120 \times 2} = \frac{5040}{240} = 21$$

\]

Step 2: Calculate (0.8^5)

$$0.8^5 = 0.8 \times 0.8 \times 0.8 \times 0.8 \times 0.8$$

Calculations:

- $(0.8 \times 0.8 = 0.64)$
- $(0.64 \times 0.8 = 0.512)$
- $(0.512 \times 0.8 = 0.4096)$
- $(0.4096 \times 0.8 = 0.32768)$

Thus:

$$0.8^5 = 0.32768$$

Step 3: Calculate (0.2^2)

$$0.2^2 = 0.2 \times 0.2 = 0.04$$

Step 4: Combine all components

Putting everything together:

$$P(x=5) = 21 \times 0.32768 \times 0.04$$

Multiply (0.32768×0.04) :

$$0.32768 \times 0.04 = 0.0131072$$

Finally, multiply by 21:

$$0.0131072 \times 21 = 0.2752512$$

$$21 \times 0.0131072 = 0.2752528$$

\]

Final Answer Rounded to Four Decimal Places

\[

$$P(x=5) \approx \boxed{0.2753}$$

\]

This means there is approximately a 27.53% chance of observing exactly 5 successes out of 7 trials when the success probability per trial is 0.8.

Interpreting the Result

Understanding how to interpret this probability is crucial for applying the binomial distribution effectively. In practical terms:

- If you repeated an experiment with 7 trials, each having an 80% success rate, you can expect to achieve exactly 5 successes roughly 27.53% of the time.
- This probability helps in risk assessment, quality control, and decision-making processes where outcomes follow a binomial pattern.
- By adjusting the parameters (N) and (P) , you can model different scenarios and compute precise probabilities for various outcomes.

Additional Insights on Binomial Probabilities

- Most Probable Number of Successes: For large (N) , the most probable number of successes is approximately $(N \times P)$. Here, $(7 \times 0.8 = 5.6)$, so 6 successes are slightly more probable than 5, but calculating specific probabilities like $(P(x=5))$ provides more detailed insights.
- Symmetry: The binomial distribution is symmetric when $(P = 0.5)$. When $(P \neq 0.5)$, it skews towards either the lower or higher number of successes.
- Use of Technology: Calculators, statistical software, or programming languages like Python or R can efficiently compute binomial probabilities, especially for larger (N) .

Conclusion

Calculating the probability $P(x=5)$ using the binomial distribution involves understanding the formula, performing factorial and exponential calculations, and carefully combining all components. For $N=7$ and $P=0.8$, the probability of exactly five successes is approximately 0.2753 when rounded to four decimal places. Mastering these calculations enhances your ability to analyze binomial scenarios accurately, informing better decision-making in various fields such as statistics, quality control, and data science.

References and Tools

- Binomial Coefficient Calculator: Online tools or programming functions (e.g., `comb` in Python's `math` module)
- Statistical Software: R (`dbinom` function), Python (`scipy.stats.binom.pmf`)
- Further Reading: "Statistics for Business and Economics" by Paul Newbold, William L. Carlson, and Betty Thorne

Understanding and applying the binomial distribution is a foundational skill in probability and statistics, enabling precise predictions and informed decisions based on discrete data.

Frequently Asked Questions

What is the probability of exactly 5 successes when $N=7$ and $P=0.8$ using the binomial distribution?

The probability $P(x=5)$ is approximately 0.2787.

How do you calculate $P(x=5)$ in a binomial distribution with $N=7$ and $P=0.8$?

Use the formula $P(x=k) = C(n, k) p^k (1-p)^{(n-k)}$. Here, $P(x=5) = C(7, 5) 0.8^5 0.2^2$, which equals $21 \cdot 0.32768 \cdot 0.04 = 0.2787$.

What is the value of the binomial coefficient $C(7,5)$?

The binomial coefficient $C(7,5)$ is 21.

How do you compute 0.8^5 and 0.2^2 for the binomial probability?

0.8^5 equals 0.32768, and 0.2^2 equals 0.04.

What is the significance of rounding the binomial probability to 4 decimals?

Rounding to 4 decimals provides a precise and standardized way to report the probability value, making it easier to compare and interpret.

If $N=7$ and $P=0.8$, what is the expected number of successes?

The expected number of successes is $N P = 7 \cdot 0.8 = 5.6$.

Can the binomial distribution be used for $N=7$ and $P=0.8$ to find probabilities for different x values?

Yes, the binomial distribution is suitable for calculating probabilities of different numbers of successes when $N=7$ and $P=0.8$.

What is the general formula for binomial probability?

$P(x=k) = C(n, k) p^k (1-p)^{(n-k)}$, where $C(n, k)$ is the binomial coefficient.

Why is the binomial distribution appropriate for this problem?

Because it models the number of successes in a fixed number of independent Bernoulli trials with the same success probability.

What software or calculator functions can be used to find $P(x=5)$ for $N=7$ and $P=0.8$?

Many calculators and software like R (`dbinom`), Excel (`BINOM.DIST`), or statistical calculators can compute binomial probabilities directly.

Additional Resources

Binomial Distribution: Calculating $P(X=5)$ When $N=7$ and $P=0.8$

Understanding the binomial distribution is fundamental in probability theory and statistics, especially when modeling the number of successes in a fixed number of independent trials with a constant probability of success. This detailed review explores the application of the binomial distribution to solve a specific problem: given $N=7$ trials and success probability $P=0.8$, what is the probability of observing exactly 5 successes? We will delve into the theoretical underpinnings, the formula derivation, step-by-step calculations, and practical implications, culminating in a precise

answer rounded to four decimal places.

Understanding the Binomial Distribution

The binomial distribution models the probability of obtaining a fixed number of successes in a fixed number of independent Bernoulli trials. Each trial has two possible outcomes: success or failure.

Key Characteristics of Binomial Distribution

- Number of trials (N): The total count of independent experiments or observations.
- Probability of success (P): The likelihood that a single trial results in success.
- Probability of failure (Q): The likelihood that a single trial results in failure, where $Q = 1 - P$.
- Number of successes (X): The random variable representing the count of successes in N trials.

Assumptions of the Binomial Model:

- Trials are independent.
- The probability of success remains constant across trials.
- Each trial results in either success or failure.

Mathematical Formula for the Binomial Distribution

The probability of observing exactly x successes in N trials is given by the binomial probability mass function (pmf):

$$P(X = x) = \binom{N}{x} \times P^x \times (1 - P)^{N - x}$$

Where:

- $\binom{N}{x}$ is the binomial coefficient, also known as "N choose x," which calculates the number of ways to select x successes from N trials.
- P^x accounts for the probability of success in the x successful trials.
- $(1 - P)^{N - x}$ accounts for the probability of failure in the remaining trials.

Binomial Coefficient Definition:

$$\binom{N}{x} = \frac{N!}{x! \times (N - x)!}$$

Where:

- $N!$ denotes factorial of N.
- $x!$ denotes factorial of x.
- $(N - x)!$ denotes factorial of $(N - x)$.

Applying the Formula to Our Problem: N=7, P=0.8, x=5

Given:

- $(N = 7)$
- $(P = 0.8)$
- $(x = 5)$

Our goal is to compute:

$$P(X=5) = \binom{7}{5} \times (0.8)^5 \times (0.2)^2$$

Step 1: Calculate the binomial coefficient $\binom{7}{5}$

$$\binom{7}{5} = \frac{7!}{5! \times (7 - 5)!} = \frac{7!}{5! \times 2!}$$

Calculate factorials:

- $7! = 7 \times 6 \times 5 \times 4 \times 3 \times 2 \times 1 = 5040$
- $5! = 5 \times 4 \times 3 \times 2 \times 1 = 120$
- $2! = 2 \times 1 = 2$

Plug in:

$$\binom{7}{5} = \frac{5040}{120 \times 2} = \frac{5040}{240} = 21$$

Step 2: Calculate $P^x = (0.8)^5$

$$(0.8)^5 = 0.8 \times 0.8 \times 0.8 \times 0.8 \times 0.8$$

Calculations:

- $0.8 \times 0.8 = 0.64$
- $0.64 \times 0.8 = 0.512$
- $0.512 \times 0.8 = 0.4096$
- $0.4096 \times 0.8 = 0.32768$

Thus,

$$(0.8)^5 \approx 0.32768$$

Step 3: Calculate $(1 - P)^{N - x} = (0.2)^2$

$$(0.2)^2 = 0.2 \times 0.2 = 0.04$$

Step 4: Combine all components

$$P(X=5) = 21 \times 0.32768 \times 0.04$$

Calculate:

- $(0.32768 \times 0.04 = 0.0131072)$
- Multiply by 21:

$$21 \times 0.0131072 = 0.2752512$$

Step 5: Round to four decimal places

$$P(X=5) \approx 0.2753$$

Interpretation of the Result

The probability of obtaining exactly 5 successes out of 7 independent trials, each with an 80% chance of success, is approximately 0.2753. This indicates that in a typical scenario, there's about a 27.53% chance of observing precisely five successes under these conditions.

This probability is relatively high, which aligns with intuition because success probability per trial (0.8) is quite high, making multiple successes within 7 trials quite probable.

Practical Applications of Binomial Probability Calculations

Understanding how to compute specific binomial probabilities has vital applications across various fields:

- Quality Control: Estimating the probability of a certain number of defective items in a batch.
- Medical Diagnostics: Calculating the likelihood of a certain number of positive test results given

test sensitivity.

- Marketing and Business: Estimating the success rate of a campaign based on response probabilities.
- Genetics: Determining the probability of inheriting a specific number of dominant alleles.
- Sports Analytics: Calculating the likelihood of a player achieving a certain number of successes (e.g., goals, hits).

In each of these applications, the core binomial formula helps quantify uncertainty and supports decision-making based on probabilistic models.

Extensions and Variations of the Binomial Distribution

While the focus here is on calculating the probability for a specific number of successes, the binomial distribution also allows for:

- Calculating cumulative probabilities: $P(X \leq x)$ or $P(X \geq x)$, useful for hypothesis testing or confidence intervals.
- Approximations: For large N , normal approximation can be used to estimate binomial probabilities, simplifying calculations.
- Binomial variance and mean: Understanding the spread and expected value:

$$\text{Mean} = N \times P$$

$$\text{Variance} = N \times P \times (1 - P)$$

For our case:

$$\text{Mean} = 7 \times 0.8 = 5.6$$

$$\text{Variance} = 7 \times 0.8 \times 0.2 = 1.12$$

The standard deviation is $\sqrt{1.12} \approx 1.0583$.

Conclusion

Calculating the probability of a specific number of successes in a binomial setting involves a clear understanding of the model's assumptions and the application of the binomial formula. In our case, with $N=7$ and $P=0.8$, computing $P(X=5)$ results in approximately 0.2753 when rounded to four decimal places.

This process exemplifies the power of the binomial distribution in probabilistic modeling, enabling analysts, statisticians, and researchers to make informed predictions and assessments in various real-world scenarios. Mastering these calculations enhances one's ability to interpret data, plan experiments, and evaluate outcomes effectively.

Summary:

- The binomial distribution provides a framework for modeling success counts.
- The probability of exactly 5 successes in 7 trials with success probability 0.8 is approximately 0.2753.
- The calculation involves factorials, binomial coefficients, and powers of success/failure probabilities.
- Practical applications span numerous fields, emphasizing the importance of understanding binomial probabilities and their calculations.

Note: For more complex or larger N , computational tools like calculator functions, statistical software (e.g., R, Python), or binomial tables are often used to facilitate rapid and accurate calculations.

[Using The Binomial Distribution If N 7 And P 0.8 Find P X 5 Round Your Answer To 4 Decimals](#)

Using The Binomial Distribution If N 7 And P 0.8 Find P X 5 Round Your Answer To 4 Decimals

Related Articles

- [B. Find The Margin Of Error. E=enter Your Response Here G \(round To One Decimal Place As Needed.\)](#)
- [Indicate Whether The Statement Is True Or False. 1. Hay Fever Is A Reaction In The Mucous Membranes](#)
- [Green Light Has A Frequency Of About 6. 001014s1. What Is The Energy Of A Photon Of Green Light?](#)

[Back to Home](#)