

# A NURSE IS PROVIDING TEACHING FOR A CLIENT WHO HAS CIRRHOSIS AND A NEW PRESCRIPTION FOR LACTULOSE. THE

A NURSE IS PROVIDING TEACHING FOR A CLIENT WHO HAS CIRRHOSIS AND A NEW PRESCRIPTION FOR LACTULOSE. THE FOLLOWING GUIDE AIMS TO DELIVER COMPREHENSIVE, SEO-OPTIMIZED INFORMATION ABOUT CIRRHOSIS MANAGEMENT, FOCUSING ON PATIENT EDUCATION REGARDING LACTULOSE THERAPY. PROPER UNDERSTANDING OF THIS MEDICATION, ITS PURPOSE, ADMINISTRATION, POTENTIAL SIDE EFFECTS, AND LIFESTYLE MODIFICATIONS IS ESSENTIAL FOR EFFECTIVE DISEASE MANAGEMENT AND IMPROVED PATIENT OUTCOMES.

---

## UNDERSTANDING CIRRHOSIS: AN OVERVIEW

### WHAT IS CIRRHOSIS?

CIRRHOSIS IS A CHRONIC LIVER DISEASE CHARACTERIZED BY PROGRESSIVE SCARRING (FIBROSIS) OF THE LIVER TISSUE. THIS SCARRING IMPAIRS NORMAL LIVER FUNCTION AND DISRUPTS BLOOD FLOW WITHIN THE LIVER, LEADING TO VARIOUS COMPLICATIONS. CIRRHOSIS CAN RESULT FROM MULTIPLE CAUSES, INCLUDING CHRONIC ALCOHOL CONSUMPTION, VIRAL HEPATITIS (B AND C), FATTY LIVER DISEASE, AUTOIMMUNE LIVER DISEASES, AND CERTAIN INHERITED DISORDERS.

### SIGNS AND SYMPTOMS OF CIRRHOSIS

PATIENTS WITH CIRRHOSIS MAY EXHIBIT A RANGE OF SYMPTOMS, SUCH AS:

- FATIGUE AND WEAKNESS
- JAUNDICE (YELLOWING OF THE SKIN AND EYES)
- EDEMA AND ASCITES (FLUID ACCUMULATION IN THE ABDOMEN)
- EASY BRUISING AND BLEEDING
- ITCHING
- LOSS OF APPETITE AND WEIGHT LOSS
- CONFUSION OR HEPATIC ENCEPHALOPATHY

### COMPLICATIONS ASSOCIATED WITH CIRRHOSIS

CIRRHOSIS CAN LEAD TO SERIOUS COMPLICATIONS, INCLUDING:

- PORTAL HYPERTENSION
- VARICEAL BLEEDING
- HEPATIC ENCEPHALOPATHY
- KIDNEY DYSFUNCTION
- INCREASED RISK OF LIVER CANCER

---

## THE ROLE OF LACTULOSE IN CIRRHOSIS MANAGEMENT

### WHAT IS LACTULOSE?

LACTULOSE IS A SYNTHETIC SUGAR (DISACCHARIDE) USED PRIMARILY TO TREAT HEPATIC ENCEPHALOPATHY, A COMPLICATION

OF CIRRHOSIS. IT WORKS BY REDUCING AMMONIA LEVELS IN THE BLOOD, WHICH IS CRITICAL BECAUSE ELEVATED AMMONIA CAN CAUSE MENTAL CONFUSION AND COMA.

## How Does Lactulose Work?

LACTULOSE FUNCTIONS THROUGH SEVERAL MECHANISMS:

- IT ACIDIFIES THE COLON CONTENTS, CONVERTING AMMONIA ( $\text{NH}_3$ ) INTO AMMONIUM ( $\text{NH}_4^+$ ), WHICH IS LESS ABSORBABLE.
- IT ACTS AS AN OSMOTIC LAXATIVE, INCREASING STOOL FREQUENCY AND VOLUME, WHICH HELPS ELIMINATE AMMONIA-PRODUCING BACTERIA AND AMMONIA ITSELF.
- IT ALTERS GUT FLORA, DECREASING THE PRODUCTION OF AMMONIA.

## INDICATIONS FOR LACTULOSE USE

LACTULOSE IS PRESCRIBED TO:

- TREAT HEPATIC ENCEPHALOPATHY
- PREVENT EPISODES OF HEPATIC ENCEPHALOPATHY IN HIGH-RISK PATIENTS
- REDUCE SERUM AMMONIA LEVELS

---

## PATIENT EDUCATION: PRESCRIBING AND ADMINISTERING LACTULOSE

### UNDERSTANDING THE PRESCRIPTION

WHEN A CLIENT WITH CIRRHOSIS RECEIVES A NEW PRESCRIPTION FOR LACTULOSE, EDUCATION SHOULD FOCUS ON:

- THE PURPOSE OF THE MEDICATION IN MANAGING HEPATIC ENCEPHALOPATHY
- HOW IT HELPS IMPROVE MENTAL STATUS AND REDUCE AMMONIA LEVELS
- THE IMPORTANCE OF ADHERING TO PRESCRIBED DOSAGE AND SCHEDULE

### How to Take Lactulose

LACTULOSE ADMINISTRATION INVOLVES:

- FOLLOWING THE SPECIFIC DOSING INSTRUCTIONS PROVIDED BY THE HEALTHCARE PROVIDER
- TYPICALLY GIVEN ORALLY, WITH DOSES ADJUSTED TO PRODUCE 2-3 SOFT STOOLS PER DAY
- USING MEASURING DEVICES TO ENSURE ACCURATE DOSING
- MAINTAINING CONSISTENT TIMING TO OPTIMIZE THERAPEUTIC EFFECTS

### MONITORING AND ADJUSTING DOSAGE

- THE GOAL IS TO ACHIEVE 2-3 SOFT, FORMED STOOLS DAILY WITHOUT DIARRHEA
- REGULAR ASSESSMENT OF STOOL PATTERN AND MENTAL STATUS IS ESSENTIAL
- DOSAGE ADJUSTMENTS SHOULD BE MADE BASED ON CLINICAL RESPONSE AND LAB RESULTS

### POTENTIAL SIDE EFFECTS AND MANAGEMENT

COMMON SIDE EFFECTS INCLUDE:

- DIARRHEA
- ABDOMINAL CRAMPING AND BLOATING
- NAUSEA
- ELECTROLYTE IMBALANCES (E.G., HYPOKALEMIA, DEHYDRATION)

PATIENTS SHOULD REPORT:

- SEVERE DIARRHEA OR DEHYDRATION SYMPTOMS
- ABDOMINAL PAIN OR CRAMPING
- SIGNS OF ELECTROLYTE IMBALANCE SUCH AS MUSCLE WEAKNESS OR IRREGULAR HEARTBEAT

---

## PRECAUTIONS AND SAFETY CONSIDERATIONS

### MONITORING LABORATORY VALUES

REGULAR LAB TESTS ARE CRUCIAL TO MONITOR:

- SERUM AMMONIA LEVELS
- ELECTROLYTES (POTASSIUM, SODIUM)
- LIVER FUNCTION TESTS

### HYDRATION AND DIETARY RECOMMENDATIONS

- ENCOURAGE ADEQUATE FLUID INTAKE UNLESS CONTRAINDICATED
- RECOMMEND A BALANCED DIET WITH SUFFICIENT PROTEIN INTAKE AS ADVISED BY THE HEALTHCARE PROVIDER
- LIMIT OR AVOID ALCOHOL CONSUMPTION

### PATIENT SAFETY TIPS

- STORE LACTULOSE OUT OF REACH OF CHILDREN
- USE THE CORRECT MEASURING DEVICE
- DO NOT DISCONTINUE MEDICATION WITHOUT MEDICAL ADVICE
- KEEP FOLLOW-UP APPOINTMENTS FOR ONGOING ASSESSMENT

---

## LIFESTYLE MODIFICATIONS FOR CLIENTS WITH CIRRHOSIS

### DIETARY GUIDELINES

- CONSUME A HIGH-PROTEIN, LOW-SODIUM DIET TO PREVENT ASCITES
- LIMIT FOODS HIGH IN SALT
- AVOID RAW OR UNDERCOOKED SHELLFISH TO REDUCE INFECTION RISK
- FOLLOW SPECIFIC DIETARY RECOMMENDATIONS PROVIDED BY A HEALTHCARE PROFESSIONAL

### ALCOHOL AND SUBSTANCE USE

- COMPLETE ABSTINENCE FROM ALCOHOL IS MANDATORY
- SEEK SUPPORT PROGRAMS FOR ALCOHOL DEPENDENCE IF NEEDED

## MANAGING COMPLICATIONS

- REGULARLY MONITOR FOR SIGNS OF BLEEDING, INFECTION, OR WORSENING MENTAL STATUS
- REPORT ANY SUDDEN CHANGES IN SYMPTOMS PROMPTLY

## IMPORTANCE OF FOLLOW-UP AND COMMUNICATION

EFFECTIVE MANAGEMENT OF CIRRHOSIS INVOLVES:

- REGULAR MEDICAL FOLLOW-UP APPOINTMENTS
- BLOOD TESTS TO MONITOR DISEASE PROGRESSION
- OPEN COMMUNICATION WITH HEALTHCARE PROVIDERS REGARDING MEDICATION EFFECTS AND SIDE EFFECTS

---

## SUMMARY: KEY POINTS FOR PATIENTS PRESCRIBED LACTULOSE

- UNDERSTAND THE PURPOSE OF LACTULOSE IN MANAGING HEPATIC ENCEPHALOPATHY
- FOLLOW DOSING INSTRUCTIONS CAREFULLY TO MAINTAIN EFFECTIVE STOOL CONSISTENCY
- BE AWARE OF POTENTIAL SIDE EFFECTS AND WHEN TO SEEK MEDICAL ATTENTION
- MAINTAIN A LIVER-FRIENDLY DIET AND ABSTAIN FROM ALCOHOL
- KEEP ALL FOLLOW-UP APPOINTMENTS AND LAB TESTS FOR OPTIMAL MANAGEMENT
- COMMUNICATE OPENLY WITH HEALTHCARE PROVIDERS ABOUT ANY CONCERNS OR SYMPTOMS

---

## CONCLUSION

PATIENT EDUCATION IS A CORNERSTONE OF EFFECTIVE CIRRHOSIS MANAGEMENT, PARTICULARLY WHEN INITIATING THERAPIES LIKE LACTULOSE. A NURSE PLAYS A VITAL ROLE IN ENSURING CLIENTS UNDERSTAND THEIR MEDICATION REGIMEN, LIFESTYLE MODIFICATIONS, AND THE IMPORTANCE OF MONITORING. THROUGH COMPREHENSIVE TEACHING AND SUPPORT, CLIENTS CAN BETTER MANAGE THEIR CONDITION, MINIMIZE COMPLICATIONS, AND IMPROVE QUALITY OF LIFE.

---

KEYWORDS FOR SEO OPTIMIZATION:

- CIRRHOSIS MANAGEMENT
- LACTULOSE FOR HEPATIC ENCEPHALOPATHY
- LIVER DISEASE PATIENT EDUCATION
- MANAGING CIRRHOSIS COMPLICATIONS
- LACTULOSE DOSAGE AND SIDE EFFECTS
- CIRRHOSIS LIFESTYLE MODIFICATIONS
- HEPATIC ENCEPHALOPATHY TREATMENT
- NURSE TEACHING FOR LIVER DISEASE

## FREQUENTLY ASKED QUESTIONS

### WHAT IS THE PRIMARY PURPOSE OF PRESCRIBING LACTULOSE FOR A CLIENT WITH CIRRHOSIS?

LACTULOSE IS PRESCRIBED TO REDUCE ELEVATED AMMONIA LEVELS IN THE BLOOD BY DECREASING AMMONIA ABSORPTION FROM THE INTESTINES, THEREBY PREVENTING HEPATIC ENCEPHALOPATHY IN CLIENTS WITH CIRRHOSIS.

## How should I take Lactulose to ensure effective treatment?

Take Lactulose exactly as prescribed, usually 2-3 soft stools per day, and adjust the dose as directed by your healthcare provider to maintain this effect. It's important not to take more or less without consulting your provider.

## What are common side effects of Lactulose that I should watch for?

Common side effects include diarrhea, bloating, abdominal cramps, and nausea. If diarrhea becomes severe or persistent, contact your healthcare provider promptly.

## Are there any dietary restrictions I should follow while taking Lactulose?

While taking Lactulose, it's important to maintain a balanced diet. Your healthcare provider may recommend limiting protein intake if ammonia levels are high, but always follow their specific dietary guidance.

## How can I tell if Lactulose is effectively reducing my ammonia levels?

Effectiveness is usually monitored through blood tests measuring ammonia levels. Additionally, a reduction in symptoms of hepatic encephalopathy, such as confusion or forgetfulness, indicates improvement.

## What should I do if I experience severe diarrhea or dehydration while on Lactulose?

If you experience severe diarrhea, dehydration, or signs of electrolyte imbalance, contact your healthcare provider immediately. They may need to adjust your dose or provide additional treatment.

## Are there any medications or substances I should avoid while taking Lactulose?

Avoid excessive use of laxatives or antidiarrheal medications without medical advice, as they can interfere with Lactulose's effectiveness or cause adverse effects. Always inform your healthcare provider of all medications and supplements you are taking.

## Additional Resources

A Nurse Is Providing Teaching For A Client Who Has Cirrhosis And A New Prescription For Lactulose

When managing chronic liver conditions such as cirrhosis, healthcare providers must navigate a complex web of treatment strategies aimed at mitigating symptoms, preventing complications, and improving quality of life. Among the medications prescribed for patients with cirrhosis, Lactulose plays a crucial role, especially in managing hepatic encephalopathy—a common and serious complication of liver failure. As a nurse, providing comprehensive patient education on the use of Lactulose is essential to ensure safety, adherence, and optimal outcomes. This guide offers a detailed overview of how to educate clients who have cirrhosis and have been prescribed Lactulose, emphasizing understanding, administration, side effects, and lifestyle considerations.

---

Understanding Cirrhosis and the Role of Lactulose

What Is Cirrhosis?

Cirrhosis is a chronic liver disease characterized by irreversible scarring (fibrosis) of liver tissue. It results from prolonged liver injury caused by factors such as chronic alcohol abuse, hepatitis B or C infections, non-

ALCOHOLIC FATTY LIVER DISEASE, OR OTHER LIVER INSULTS. OVER TIME, CIRRHOSIS IMPAIRS THE LIVER'S ABILITY TO PERFORM VITAL FUNCTIONS, INCLUDING DETOXIFICATION, SYNTHESIS OF PROTEINS, AND PRODUCTION OF CLOTTING FACTORS.

## WHY IS LACTULOSE PRESCRIBED?

LACTULOSE IS A SYNTHETIC SUGAR USED PRIMARILY TO TREAT HEPATIC ENCEPHALOPATHY—A DECLINE IN BRAIN FUNCTION RESULTING FROM THE LIVER'S INABILITY TO DETOXYFY AMMONIA AND OTHER NEUROTOXINS. ELEVATED AMMONIA LEVELS CAN CROSS THE BLOOD-BRAIN BARRIER, LEADING TO CONFUSION, ALTERED MENTAL STATUS, AND EVEN COMA.

LACTULOSE WORKS BY:

- REDUCING AMMONIA ABSORPTION IN THE INTESTINES
- PROMOTING EXCRETION OF AMMONIA THROUGH INCREASED STOOL FREQUENCY
- ALTERING GUT FLORA TO DECREASE AMMONIA-PRODUCING BACTERIA

---

## KEY PRINCIPLES OF TEACHING ABOUT LACTULOSE

PROVIDING EFFECTIVE EDUCATION INVOLVES ADDRESSING SEVERAL CORE AREAS:

- UNDERSTANDING THE MEDICATION'S PURPOSE
- PROPER ADMINISTRATION TECHNIQUES
- RECOGNIZING AND MANAGING SIDE EFFECTS
- MONITORING AND FOLLOW-UP
- LIFESTYLE AND DIETARY CONSIDERATIONS

---

## EXPLAINING THE PURPOSE AND MECHANISM OF LACTULOSE

### CLARIFY THE MEDICATION'S ROLE

BEGIN BY EXPLAINING THAT LACTULOSE IS PRESCRIBED TO HELP REDUCE THE BUILDUP OF TOXINS LIKE AMMONIA IN THE BODY, WHICH CAN CAUSE CONFUSION OR OTHER MENTAL CHANGES IN CIRRHOSIS PATIENTS. EMPHASIZE THAT LACTULOSE IS A PREVENTIVE AND THERAPEUTIC MEDICATION AIMED AT MAINTAINING BRAIN FUNCTION AND PREVENTING EPISODES OF HEPATIC ENCEPHALOPATHY.

### HOW DOES IT WORK?

EXPLAIN THAT LACTULOSE:

- IS A SYNTHETIC SUGAR THAT IS NOT ABSORBED INTO THE BLOODSTREAM
- PASSES THROUGH THE DIGESTIVE SYSTEM UNCHANGED
- IS FERMENTED BY BACTERIA IN THE GUT, PRODUCING ACIDS THAT CONVERT AMMONIA INTO A FORM THAT CAN BE EXPELLED IN STOOL
- ACTS AS A LAXATIVE TO HELP CLEAR AMMONIA FROM THE INTESTINES

---

## PROPER ADMINISTRATION AND DOSAGE

### TYPICAL DOSAGE AND SCHEDULE

LACTULOSE DOSAGE VARIES BASED ON INDIVIDUAL NEEDS, BUT COMMON STARTING POINTS INCLUDE:

- INITIAL DOSE: 15-30 mL (OR 45-60 GRAMS) ORALLY, 2-3 TIMES DAILY
- ADJUSTMENTS: BASED ON STOOL FREQUENCY AND AMMONIA LEVELS, OFTEN AIMING FOR 2-3 SOFT STOOLS PER DAY

### ADMINISTRATION TIPS

- TIMING: TAKE LACTULOSE WITH OR WITHOUT FOOD. CONSISTENCY IN TIMING HELPS MAINTAIN STABLE STOOL PATTERNS.
- MEASURING: USE AN ACCURATE MEASURING DEVICE OR CUP FOR LIQUID FORMULATIONS.

- MIXING: IT CAN BE MIXED WITH WATER, JUICE, OR OTHER FLUIDS IF THE CLIENT PREFERS, MAKING IT EASIER TO SWALLOW.
- STOOL MONITORING: ENCOURAGE CLIENTS TO TRACK THEIR BOWEL MOVEMENTS TO HELP ADJUST DOSAGE APPROPRIATELY.

## ADHERENCE STRATEGIES

- REINFORCE THE IMPORTANCE OF TAKING THE MEDICATION EXACTLY AS PRESCRIBED.
- DISCUSS BARRIERS SUCH AS TASTE, FREQUENCY, OR SIDE EFFECTS THAT MAY AFFECT ADHERENCE.
- PROVIDE TIPS FOR REMEMBERING DOSES, SUCH AS SETTING ALARMS OR INCORPORATING MEDICATION INTO DAILY ROUTINES.

---

## RECOGNIZING AND MANAGING SIDE EFFECTS

### COMMON SIDE EFFECTS

CLIENTS SHOULD BE AWARE OF POTENTIAL ADVERSE EFFECTS, INCLUDING:

- DIARRHEA: EXCESSIVE BOWEL MOVEMENTS LEADING TO DEHYDRATION
- ABDOMINAL CRAMPING OR BLOATING
- NAUSEA
- ELECTROLYTE IMBALANCES (E.G., HYPOKALEMIA, HYPONATREMIA)

### WHEN TO SEEK MEDICAL ATTENTION

PATIENTS SHOULD BE INSTRUCTED TO REPORT:

- SEVERE DIARRHEA OR DEHYDRATION SIGNS
- ABDOMINAL PAIN OR CRAMPING
- PERSISTENT NAUSEA OR VOMITING
- SYMPTOMS OF ELECTROLYTE IMBALANCE (WEAKNESS, IRREGULAR HEARTBEAT)
- ANY SIGNS OF ALLERGIC REACTION (RASH, SWELLING, DIFFICULTY BREATHING)

### MANAGING SIDE EFFECTS

- ADJUST DOSAGE UNDER MEDICAL SUPERVISION
- MAINTAIN ADEQUATE HYDRATION
- MONITOR ELECTROLYTE LEVELS REGULARLY

---

## MONITORING AND FOLLOW-UP

### LABORATORY TESTS

ROUTINE MONITORING INCLUDES:

- SERUM AMMONIA LEVELS: TO EVALUATE EFFECTIVENESS
- ELECTROLYTES: ESPECIALLY POTASSIUM AND SODIUM
- LIVER FUNCTION TESTS: TO ASSESS DISEASE PROGRESSION

### CLINICAL ASSESSMENTS

- OBSERVE MENTAL STATUS FOR SIGNS OF IMPROVEMENT OR DETERIORATION
- MONITOR BOWEL MOVEMENT FREQUENCY AND STOOL CONSISTENCY
- ASSESS HYDRATION STATUS AND ELECTROLYTE BALANCE

### PATIENT COMMUNICATION

ENCOURAGE CLIENTS TO:

- KEEP A SYMPTOM DIARY
- REPORT ANY CONCERNS OR WORSENING SYMPTOMS PROMPTLY
- ATTEND REGULAR FOLLOW-UP APPOINTMENTS

---

## DIETARY AND LIFESTYLE CONSIDERATIONS

### DIETARY RECOMMENDATIONS

WHILE LACTULOSE HELPS REDUCE AMMONIA LEVELS, DIET ALSO PLAYS A VITAL ROLE:

- PROTEIN INTAKE: MODERATE, CONSISTENT INTAKE TO PREVENT MALNUTRITION BUT AVOID EXCESS AMMONIA PRODUCTION
- AVOIDANCE OF HIGH-PROTEIN FOODS: SUCH AS CERTAIN MEATS, DAIRY, OR BEANS IF ADVISED BY THE HEALTHCARE TEAM
- HYDRATION: ADEQUATE FLUID INTAKE TO PREVENT DEHYDRATION AND SUPPORT STOOL SOFTENING

### LIFESTYLE MODIFICATIONS

- ABSTAIN FROM ALCOHOL
- AVOID MEDICATIONS OR SUBSTANCES THAT CAN IMPAIR LIVER FUNCTION OR CAUSE GASTROINTESTINAL IRRITATION
- PRACTICE GOOD HAND HYGIENE TO PREVENT INFECTIONS, ESPECIALLY IN IMMUNOCOMPROMISED PATIENTS

---

## PATIENT EDUCATION AND ENGAGEMENT

### EMPOWERING THE CLIENT

- EXPLAIN THE IMPORTANCE OF MEDICATION ADHERENCE
- CLARIFY MISCONCEPTIONS (E.G., LACTULOSE IS NOT A CURE BUT A MANAGEMENT TOOL)
- REINFORCE THAT SIDE EFFECTS ARE MANAGEABLE AND SHOULD BE COMMUNICATED TO HEALTHCARE PROVIDERS
- ENCOURAGE QUESTIONS AND ACTIVE PARTICIPATION IN CARE PLANNING

### SUPPORT SYSTEMS

- INVOLVE FAMILY OR CAREGIVERS IN EDUCATION
- PROVIDE WRITTEN INSTRUCTIONS OR EDUCATIONAL MATERIALS
- CONNECT CLIENTS WITH SUPPORT GROUPS OR COUNSELING SERVICES AS NEEDED

---

## CONCLUSION

PROVIDING THOROUGH, PATIENT-CENTERED EDUCATION ABOUT LACTULOSE FOR CLIENTS WITH CIRRHOSIS IS ESSENTIAL IN MANAGING HEPATIC ENCEPHALOPATHY EFFECTIVELY. NURSES PLAY A VITAL ROLE IN ENSURING CLIENTS UNDERSTAND HOW TO TAKE THEIR MEDICATION CORRECTLY, RECOGNIZE SIDE EFFECTS, AND MAKE APPROPRIATE LIFESTYLE ADJUSTMENTS. THROUGH CLEAR COMMUNICATION, DILIGENT MONITORING, AND ONGOING SUPPORT, HEALTHCARE PROVIDERS CAN HELP IMPROVE TREATMENT ADHERENCE, PREVENT COMPLICATIONS, AND ENHANCE THE OVERALL QUALITY OF LIFE FOR INDIVIDUALS LIVING WITH CIRRHOSIS.

---

REMEMBER: EACH PATIENT IS UNIQUE, AND EDUCATION SHOULD BE TAILORED TO INDIVIDUAL NEEDS, LITERACY LEVELS, AND CULTURAL CONSIDERATIONS. ALWAYS COLLABORATE WITH THE HEALTHCARE TEAM TO OPTIMIZE CARE PLANS AND ENSURE SAFETY.

## **A Nurse Is Providing Teaching For A Client Who Has Cirrhosis And A New Prescription For Lactulose The**

A Nurse Is Providing Teaching For A Client Who Has Cirrhosis And A New Prescription For Lactulose The

## Related Articles

- [A Student Has A Solution Of Unknown Concentration Of The Strong Acid HCl. Which Of The Following Would](#)
- [A Number Of Factors Influence The Sample Size For A Substantive Test Of Details Of An Account Balance.](#)
- [A Loop With An Ac Current Produces A Changing Magnetic Field. Two Loops Have The Same Area, But One Is](#)

[Back to Home](#)